International Revista of Physics, Chemistry, Mathematics and Biology
(ISSN: xxxx-xxxx)
Vol1, Issue 1, (2026)
https://irpcmb.com/

Bioelectromagnetic Responses of Cellular Systems

Dr. William George Parker

Research Fellow, Centre for Entrepreneurship and Small Business Development, Nordic Business Institute, Copenhagen, Denmark.

Citation: Dr. William George Parker (2026) Bioelectromagnetic Responses of Cellular Systems , *International Revista of Physics, Chemistry, Mathematics and Biology, Vol. 1: Issue 1 (2026).*

Received: Mar 20, 2026; **Accepted**: May 10, 2026; **Published**: June 28, 2026

Abstract

Cellular bioelectromagnetics examines how defined electric and magnetic exposures influence cells. Studies have reported changes in intracellular calcium, redox balance, gene expression, proliferation and differentiation under selected conditions. The findings depend strongly on frequency, field strength, waveform, duration, cell identity and culture environment. This narrative review distinguishes static and extremely low frequency fields from radiofrequency exposure; summarizes proposed signaling pathways; and discusses experimental evidence from neuronal and bone-cell models. A two-study chart depicts exposure settings rather than comparing biological effect sizes. Cellular responses in vitro establish neither a universal mechanism nor a human health outcome. Reproducible dosimetry, sham exposure, temperature control and independent replication are essential for interpreting and translating results.

Keywords:bioelectromagnetics; electromagnetic fields; calcium signaling; oxidative stress; pulsed electromagnetic fields; cellular response

1 Introduction

Living cells maintain membrane voltage, ionic gradients and redox balance. External electromagnetic fields can interact with biological tissue through different physical routes. At extremely low frequencies (ELF), time-varying magnetic fields induce electric fields; sufficiently strong exposures stimulate nerves and muscles. At radiofrequency (RF), absorbed energy can heat tissue. Proposed responses to weaker exposures are more context dependent, and an observed molecular change is not itself proof of harm or clinical benefit [1,2].

This review focuses on controlled cell experiments and their interpretation. It includes no original biological experiments or pooled effect-size estimates. The goal is to explain which exposure variables must be reported, what cellular endpoints have been investigated and why results from different devices are not automatically comparable.

2 Exposure classes and experimental design

Static magnetic fields, ELF fields and RF fields differ in frequency and coupling to tissues. A laboratory report should specify electric-field strength or magnetic flux density at the sample, frequency, waveform, duty cycle, orientation, exposure duration, medium, container geometry and temperature. Nominal coil settings are not a substitute for measured sample-level fields. ELF cellular studies span diverse combinations of intensity, frequency, time and continuous or intermittent exposure; a scoping review classified 79 eligible articles across these dimensions [3].

Copyright © 2026 by Author/s and Licensed by IRPCMB, India. This is an open access article distributed under the Creative Common Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Conceptual sequence of candidate cellular responses. Each arrow requires direct experimental validation.

A rigorous design uses a physically comparable sham condition with the same incubator handling, vibration, coil heat and timing. Cell passage number, density and medium composition should match. Temperature probes and blinded image or assay analysis reduce bias. Investigators should predefine a primary endpoint, adequate biological replication and an analysis plan; running many biomarkers without correction increases the chance of false positive findings.

3 Proposed mechanisms

Calcium signaling is frequently investigated because channel activity and intracellular stores influence cell fate. EMF-associated calcium changes have been discussed in stem-cell models, including osteogenic and chondrogenic differentiation [4]. Mitochondrial and enzymatic redox pathways are another candidate route. Reactive oxygen species (ROS) serve as normal signals at some concentrations and can contribute to damage at others; the direction and meaning of change depend on context [7].

Heat shock proteins and stress-response genes have also been reported after some exposures [8]. Mechanistic claims need more than a changed biomarker: blocking the proposed channel, rescuing a phenotype, measuring exposure inside the culture vessel and excluding heating are stronger tests of causality. A response observed in one cell line at one field setting should not be generalized to unrelated cells.

4 Cellular endpoints and representative evidence

Table 1. Selected published cellular models. Conditions and endpoints differ, so rows are not effect-size comparisons.

Cell system	Field example	Reported endpoint and interpretation
Human SH-SY5Y neuronal cells	50 Hz, 1 mT ELF magnetic exposure [6]	Oxidative stress and neuronal-cell responses studied; culture result does not establish neurological disease risk.
Irradiated osteoblasts	15 Hz, 2 mT pulsed-burst EMF [5]	Distinct intracellular calcium oscillations and bone-cell protection investigated in an experimental setting.
Stem cells	Varying ELF or pulsed conditions [4]	Calcium-associated differentiation and cell-fate pathways discussed across heterogeneous studies.
Mouse preosteoblast MC3T3-E1	A clinically used PEMF signal [9]	Osteoblastic differentiation with adenosine-receptor-related gene activation reported.

The two numerical exposure settings in Table 1 are illustrated below. Frequency and magnetic flux density are properties of the applied field, not measures of biological response. The studies used different cells, waveforms and research questions; their locations on the chart cannot rank effectiveness or safety.




Figure 2. Published settings from an osteoblast experiment [5] and a neuronal-cell experiment [6]. No outcome data are plotted.

5 Interpretation and translational boundaries

A field can yield different outcomes in proliferating versus differentiated cells and in healthy versus stressed cultures. Nonlinear patterns have been proposed, including possible frequency or amplitude windows, but mechanisms remain debated [7]. A null result at one exposure does not refute every possible effect; a positive result does not validate every frequency. The relevant inference is restricted to the measured cell type, field configuration and endpoint.

Bone repair is an active application area for pulsed electromagnetic fields. One experimental study identified a 15 Hz, 2 mT pulsed-burst setting associated with calcium spikes in irradiated osteoblasts [5]. Other preosteoblast work reported adenosine receptor involvement in differentiation [9]. Those mechanistic observations warrant controlled translational testing. They should not be presented as direct evidence that an arbitrary consumer EMF device treats bone disease.

For public-health questions, exposure assessment and disease endpoints require population and animal evidence in addition to cell assays. WHO distinguishes established effects at sufficiently high ELF field strengths, such as induced nerve and muscle stimulation, from unresolved questions about prolonged lower-level exposure [1]. WHO also notes that laboratory cellular changes do not automatically demonstrate a causal relationship to disease [2].

6 Research priorities

Future reports should publish field maps and waveform traces at the sample, quantify background fields and incubator temperature, and disclose coil calibration and sham procedures. Standardized dose descriptors would enable replication. Multi-laboratory studies should test prespecified conditions across several cell types with both positive and negative controls. Parallel measurement of calcium, ROS, viability and phenotype over time could distinguish transient adaptation from sustained injury or beneficial differentiation.

For clinical translation, a sequence of independent replication, relevant tissue models, animal studies and controlled trials is needed. Biological effect, therapeutic efficacy and safety are separate questions. Exposure standards and clinical decisions should be based on the totality of evidence for a specified application, rather than a single cellular readout.

7 Conclusion

Cellular systems can show measurable responses to selected electromagnetic exposures, including calcium signaling, redox changes and differentiation. The response is conditional on exposure details and biological context. The most useful next advance is reproducible sample-level dosimetry combined with sham-controlled, mechanistic experiments. Careful separation of cellular observations from health claims will make this research more informative.

References

[1] World Health Organization. Extremely low frequency fields: environmental health criteria and exposure summary. https://www.who.int/teams/environment-climate-change-and-health/radiation-and-health/non-ionizing/exposure

[2] World Health Organization. Radiation: electromagnetic fields, questions and answers. https://www.who.int/news-room/questions-and-answers/item/radiation-electromagnetic-fields

[3] The Cellular Response Is Determined by a Combination of Different ELF-EMF Exposure Parameters: A Scope Review. (2024). PubMed PMID 38791113. https://pubmed.ncbi.nlm.nih.gov/38791113/

[4] Electromagnetic fields regulate calcium-mediated cell fate of stem cells: osteogenesis, chondrogenesis and apoptosis. (2023). Stem Cell Research & Therapy, 14, 133. https://doi.org/10.1186/s13287-023-03303-w

[5] High-specificity protection against radiation-induced bone loss by a pulsed electromagnetic field. (2022). Proceedings of the National Academy of Sciences. https://pmc.ncbi.nlm.nih.gov/articles/PMC9401628/

[6] Neuronal cellular responses to extremely low frequency electromagnetic field exposure: implications regarding oxidative stress and neurodegeneration. (2014). PLOS ONE. https://doi.org/10.1371/journal.pone.0104973

[7] Molecular Biological Effects of Weak Low-Frequency Magnetic Fields: Frequency–Amplitude Efficiency Windows and Possible Mechanisms. (2023). International Journal of Molecular Sciences. https://pmc.ncbi.nlm.nih.gov/articles/PMC10342092/

[8] Blank, M., & Goodman, R. (2009). Electromagnetic fields stress living cells. Pathophysiology. https://doi.org/10.1016/j.pathophys.2009.01.006

[9] Pulsed-electromagnetic-field induced osteoblast differentiation requires activation of genes downstream of adenosine receptors A2A and A3. (2021). PLOS ONE. https://pmc.ncbi.nlm.nih.gov/articles/PMC7906300/